

Seroepidemiology of Human Parvovirus B19 in Taiwan

Kuei-Hsiang Lin,^{1*} San-Lin You,² Chien-Jen Chen,² Chu-Fung Wang,¹ Czau-Siung Yang,³ and Shudo Yamazaki⁴

¹Department of Clinical Laboratory, Kaohsiung Medical College, Kaohsiung, Taiwan, Republic of China

²Graduate Institute of Epidemiology, College of Public Health, National Taiwan University, Taipei, Taiwan

³Graduate Institute of Microbiology, College of Medicine, National Taiwan University, Taipei, Taiwan

⁴National Institute of Infectious Diseases, Toyama, Sinjuku-ku, Tokyo, Japan

In order to determine the prevalence and risk factors of human parvovirus B19 (B19) infection in Taiwan, a seroepidemiological study was carried out in 19 townships. Serum samples were collected from 862 healthy residents, who were selected by stratified random sampling from various study areas. They were chosen from four different ethnic groups including aborigines, Fukien Taiwanese, Hakka Taiwanese, and mainland Chinese. Serum samples were screened for B19 IgG antibody by indirect antibody capture enzyme-linked immunosorbent assay (ELISA) and B19 IgM by IgM antibody capture (MAC)-ELISA, respectively. The overall prevalence of anti-B19 IgG and anti-B19 IgM was 32.8% and 0.35%, respectively. The anti-B19 seropositive rate in females was significantly higher than that of males (36.4% vs. 29.4%, $P < .001$). The age-sex-adjusted seropositive rate in urban townships (39.9%) was higher than that in aboriginal townships (30.5%, $P < .001$). The seropositive rate increased significantly with age showing a dose-response relationship ($P = 0.0001$ based on a trend test). Blood transfusion was found to be associated with an increased seropositive rate showing a multivariate-adjusted odds ratios of 1.6. *J. Med. Virol.* 57: 169–173, 1999. © 1999 Wiley-Liss, Inc.

KEY WORDS: human parvovirus B19; ELISA, capture ELISA; seroepidemiology

INTRODUCTION

Human parvovirus B19 (B19) belongs to genus *Erythrovirus* of the *Parvoviridae*, a family of small DNA viruses with single-stranded, linear genomes approximately 5 kb in length [Siegl et al., 1985]. B19 antigen was first discovered in the serum of healthy blood donors [Cossart et al., 1975]. The virus has a remark-

able tissue-tropism for erythroid progenitor cells in human bone marrow [Brown et al., 1993]. Hence, the virus is associated with a wide range of clinical manifestations such as erythema infectiosum (EI) in children [Anderson et al., 1983], acute arthritis in adults [Reid et al., 1985; White et al., 1985], and aplastic crisis in patients with chronic hemolytic anemia [Serjeant et al., 1981; Rao et al., 1983; Shade et al., 1986]. B19 may also cause chronic infection in immunocompromised patients leading to persistent anemia [Pattison et al., 1981]. The virus has also been implicated as a cause of fetal hydrops and fetal loss during pregnancy [Brown et al., 1984]. Moreover, B19 may spread through the transfusion of contaminated blood products [Williams et al., 1990].

Serosurveys for B19 antibodies have been reported for Fukuoka, Japan [Nunoue et al., 1985], U.S.A. [Anderson et al., 1986], West Germany [Schwarz et al., 1987], England and Wales [Cohen and Buckley, 1988], and Hong Kong [Lim et al., 1997]. Antibody prevalence ranged from 20% to 80%, depending on the simultaneous occurrence of B19 outbreaks. The prevalence of B19 virus infection and its associated risk factors in Taiwan have never been reported. To estimate the prevalence of B19 virus infection in Taiwan, serum samples collected from 862 healthy subjects recruited in 1989 were tested for the presence of B19 virus IgG and IgM antibodies.

MATERIALS AND METHODS

Study Townships and Subjects

The selection of study townships and subjects have been described in detail previously [Wang et al., 1988]. In brief, the general population in Taiwan was first stratified according to ethnic characteristics (i.e., aborigines, Fukien Taiwanese, Hakka Taiwanese, and

*Correspondence to: Kuei-Hsiang Lin, Department of Clinical Laboratory, Kaohsiung Medical College, 100 Shih-Chuan First Road, Kaohsiung City 807, Taiwan, Republic of China. E-mail: kuhsl@ms3.hinet.net

Accepted 1 September 1998

mainland Chinese) and residential areas (i.e., northern, southern, eastern and western islands, and offshore islets). Because there were no Hakka Taiwanese living in the offshore islets, there were only 19 strata in this study (Ministry of the Interior of R.O.C., 1985). One residential township was randomly chosen from each stratum. Study subjects were selected from the name lists updated by household registration offices using a multistage random sampling method [Wang et al., 1988]. A subset sample of 862 study subjects were selected from the original samples of 7,278 for the examination of B19 seromarkers.

Serum Collection and Questionnaire Interview

The serum samples and information regarding risk factors were collected by public health nurses in local health centers in study townships. A 10-ml blood sample was collected from each study subject and serum samples were separated and stored at -70°C until examinations. A standardized interview with a structured questionnaire was used to obtain information on risk factors including sociodemographic characteristics and disease history.

B19 IgG Antibody Detection

The B19 IgG antibody was detected by indirect antibody capture enzyme-linked immunosorbent assay (ELISA) according to the manufacturer's instructions (DENKA SEIKEN, Tokyo, Japan). Briefly, 100 μl each of standards and 1:201 diluted specimens were added to viral antigen-coated microtiter plate wells. After incubation at room temperature for 1 hr, the wells were washed and 100 μl peroxidase-labeled goat antibody against human IgG was added and incubated for another 1 hr at room temperature. After washing, 100 μl substrate solution (3,3',5,5'-tetramethyl benzidine) was added and the color was developed for 30 min at room temperature. After adding 100 μl stop reagent (0.6 N sulfuric acid) the optical density was read within 30 min at the dual filter wave length of 450/630 nm with an automatic plate reader. Cut-off values for a positive test were calculated as described by the manufacturer.

B19 IgM Antibody Detection

The B19 IgM antibody was detected by MAC-ELISA according to the protocols of the manufacturer (DENKA SEIKEN, Tokyo, Japan). Briefly, 100 μl each of standards and 1:201 diluted specimens were added to antihuman IgM antibody-coated microtiter plate wells. After a 1-hr incubation at room temperature, the wells were washed and 100 μl viral antigen was added and incubated at room temperature for 1 hr. After washing, 100 μl peroxidase-labeled goat antibody against human B19 was added and incubated for another 1 hr at room temperature. The successive procedures were the same as those in B19 IgG antibody detection.

Statistical Analysis

Odds ratios (ORs) and 95% confidence intervals (CIs) were calculated by unconditional logistic regression to

examine the association between B19 infection and potential risk factors after adjustment for potential confounding factors including sex, age, and residential area [Breslow and Day, 1992]. The *P* value for trend was calculated as level of significance of regression coefficient (indicative of amount of increase in risk per unit change in prevalence) for each ordinal variable in logistic regression model with relevant adjusting variables.

RESULTS

Seropositive Rate of B19 Antibody by Sociodemographic Characteristics

Two hundred eighty-one of the 862 study subjects were seropositive for B19 IgG antibody and negative for IgM; three male subjects were seropositive for B19 IgM, (one of whom was negative for IgG and positive for IgM and two of whom were positive for both IgG and IgM). The seropositive rates of both IgG and IgM for males and females were 29.6% (130/439) and 36.4% (154/423), respectively (Table I). Thus the total positive rate was 32.9%. The overall B19 seropositive rates in aboriginal, rural, and urban townships were 30.5%, 28.5%, and 39.9%, respectively. In all residential areas, females had a higher seropositive rate than males, and both males and females in urban townships had the highest seropositive rates. The seropositive rate in females remained significantly higher than that of males ($P < .05$) after adjustment for age and residential area. The seropositive rates were found to increase significantly with age and ranged from 10.3% in ≤ 14 -year-old group to 51.3% in ≥ 60 -year-old group. The older the subject, the higher the seropositive rate after adjustment for sex and residential area ($P < .001$). Residents in urban townships had a significantly higher seropositive rate than those in aboriginal townships ($P < .05$) after adjustment for age and sex. All study subjects were classified into three groups according to their educational levels, that is, senior high or above (203 subjects), junior high (169 subjects), and elementary and illiteracy (490 subjects) groups. The results showed that there was no significant difference in seropositive rate at different educational levels.

B19 Antibody Seropositive Rates by Blood Transfusion and Tattooing

As shown in Table II, those subjects with blood transfusion histories had a higher seropositive rate (53.6%) than those without (31.9%), after adjustment for age, sex, and residential area. There was no significant association between seropositivity of B19 and tattooing history.

Multiple Regression Analysis of Risk Factors Associated With Seropositivity of B19 Antibody

In the multiple logistic regression analysis, age, sex, residential area, and blood transfusion history were included in the regression model. As shown in Table III, age, sex, and residential area remained significantly associated with seropositivity of B19 antibody.

TABLE I. Associations With Seropositivity of Antibody Against Human Parvovirus B19 for Demographic Characteristics in Taiwan, 1989

Variable	Group	No. tested	No. positive (%)	OR (95% CI)	Adjusted OR (95% CI)
Sex	Male	439	130 (29.6) ^d	1.0 (referent)	1.0 ^a (referent)
	Female	423	154 (36.4)	1.4* (1.0–1.8)	1.4* (1.0–1.9)
Age (years)	≤14	116	12 (10.3)	1.0 (referent)	1.0 ^b (referent)
	15–19	126	20 (15.9)	1.6 (0.8–3.5)	1.6 (0.8–3.5)
	20–29	115	27 (23.5)	2.7** (1.3–5.6)	2.6* (1.3–5.5)
	30–39	136	49 (36.0)	4.9*** (2.4–9.8)	4.8*** (2.4–9.7)
	40–49	132	63 (47.7)	7.9*** (4.0–15.7)	7.8*** (3.9–15.5)
	50–59	120	53 (44.2)	6.9*** (3.4–13.8)	6.6*** (3.3–13.3)
	≥60	117	60 (51.3)	9.1*** (4.5–18.4)	9.1*** (4.5–18.5)
Residential township	Trend test			$P < .001$	$P < .001$
	Aboriginal	295	90 (30.5)	1.0 (referent)	1.0 ^c (referent)
	Rural	284	81 (28.5)	0.9 (0.6–1.3)	0.9 (0.6–1.3)
	Urban	283	113 (39.9)	1.5* (1.1–2.1)	1.5* (1.1–2.2)

OR, odds ratio; CI, confidence interval.

^aAdjusted variables included: age, residential township.^bAdjusted variables included: sex, residential township.^cAdjusted variables included: sex, age.^dInclude either IgG positive or IgM positive cases.* $P < .05$; ** $P < .01$; *** $P < .001$.

TABLE II. Associations With Seropositivity of Antibody Against Human Parvovirus B19 for History of Blood Transfusion and Tattooing in Taiwan, 1989

Variable (95% CI)	Group	No. tested	No. positive (%)	OR (95% CI)	Adjusted OR ^a
Blood transfusion	No	808	258 (31.9)	1.0 (referent)	1.0 ^a (referent)
	Yes	54	26 (53.6)	2.0* (1.1–3.4)	1.7** (1.0–3.1)
Tattoo	No	810	264 (32.6)	1.0 (referent)	1.0 (referent)
	Yes	52	20 (38.5)	1.3 (0.7–2.3)	1.1 (0.6–2.1)

OR, odds ratio; CI, confidence interval.

^aAdjusted variables included: sex, age, and residential status.* $P < .05$; ** $P < .10$.

TABLE III. Multivariate-Adjusted Odds Ratios for Various Risk Factors Associated With Seropositivity of Antibody Against Human Parvovirus B19 in Taiwan, 1989

Variable	Group	Multivariate-adjusted odds ratio (95% CI)
Sex	Male	1.0 (referent)
	Female	1.4* (1.0–1.9)
Age (yr)	≤14	1.0 (referent)
	15–19	1.6 (0.7–3.5)
	20–29	2.6* (1.2–5.5)
	30–39	4.7** (2.3–9.4)
	40–49	7.4** (3.7–14.8)
	50–59	6.4** (3.3–13.5)
	≥60	8.9** (4.4–18.0)
Residential status	Aboriginal townships	1.0 (referent)
	Rural townships	0.9 (0.6–1.4)
	Urban townships	1.5* (1.1–2.2)
Blood transfusion history	No	1.0 (referent)
	Yes	1.6*** (0.9–2.8)

CI, confidence interval.

* $P < .05$; ** $P < .01$; *** $P < .10$.

The multivariate-adjusted OR was 1.4 for females compared with males, whereas residents in urban townships had a 1.5-fold seropositivity rate compared with residents in aboriginal townships. The older the sub-

ject, the greater the multivariate-adjusted OR, which was as high as 8.9 for those older than 60 years as compared with those younger than 15 years. The multivariate-adjusted OR was 1.6 for blood transfusion history.

DISCUSSION

B19 was first identified and characterized from serum samples that were assayed for hepatitis B virus. The virus was then reported to be associated with the occurrence of aplastic crisis in patients with chronic hemolytic anemia, fifth disease, still births [Knott et al., 1984], arthritis, and persistent infection [Kurtzman et al., 1989; Anderson, 1990; Brown et al., 1994; Pattison, 1994; Musiani et al., 1995; von Pöblitzki et al., 1995; Cassinotti et al., 1997; Gray et al., 1998]. It has also been associated with the contamination of blood products [Erdman et al., 1997].

B19 infection has been reported in many countries around the world. The seropositive rate of B19 IgG antibody varies by location and time of the last B19 epidemic. In Japan, the seropositive rate was reported to be 20% in the young age groups of <20 years and >80% in the old age groups of >60 years [Yamashita et al., 1992]. In England and Wales, the seropositive rate was 5–15% in the young age groups of 1–5 years, 50–60% in older children, young adults, and women of child-bearing age, and >85% in the old age group of >70

years. In the US, the seropositive rate was 2% in young age group of <5 years and 49% in adults aged >20 years.

In Taiwan, a small outbreak of EI was observed last year (personal communication), but before 1989 no outbreaks of B19 infection had been reported. The seropositive rate of B19 antibody was found to be 10.3% in young age group of ≤ 14 years and 51.3% in old age group of ≥ 60 years. The increasing seropositive rate with age is consistent with those reported in other countries [Cossart et al., 1975; Paver and Clarke, 1976; Edwards et al., 1981; Mortimer et al., 1983; Courouce et al., 1984; Okabe et al., 1984; Anderson et al., 1986; Cohen and Buckley, 1988; Yamashita et al., 1992; Lim et al., 1997].

In this study lower prevalence rate (23.5%) was found in young adults (20–29 years). This result is similar to those observed in Japan: <20% in the 20–29-year-old group [Yamashita et al., 1992] and Hong Kong (9.5% in the 15–24-year-old group [Lim et al., 1997]), but different from those in the western countries with high prevalence rate of 49% in US [Anderson et al., 1986] and 50–60% in the UK in the young age group [Cohen and Buckley, 1988]. These results suggest that only fewer sporadic cases or asymptomatic infection of B19 might have occurred in the Far East than western countries.

The higher prevalence rate in urban areas in this study may have been because of the crowdedness of the population in the urban areas, because human parvovirus B19 is transmitted effectively after close contact exposure. This result might be attributed to the possible mode of B19 transmission by aerosol via the respiratory tract. No significant difference in B19 infection rates was observed at different educational levels.

It has been reported that B19 infection can be transmitted through parenteral routes [Musiani et al., 1995]. In this study, blood transfusion history was associated with an increased seropositive rate of B19, but no association was observed between B19 infection and tattooing history.

It is interesting that a significant difference was found in this study with a higher positive rate among females (<0.05), although no gender difference was found in Japan [Matsunaga et al., 1995]. A further study is necessary to elucidate this difference.

ACKNOWLEDGMENTS

We thank Ms. Heng-Ling Wang and Sandy Yang for preparing the samples. We are grateful to DENKA SEIKEN Company for the sponsorship of all the ELISA kits.

REFERENCES

- Anderson LJ. 1990. Human parvovirus. *Journal of Infectious Disease* 161:603–608.
- Anderson LJ, Tsou C, Parker RA, Chorba TL, Tattersall P, Mortimer PP. 1986. Detection of antibodies and antigens of human parvovirus B19 by enzyme-linked immunosorbent assay. *J Clin Microbiol* 24:522–526.
- Anderson MJ, Jones SE, Fisher-Hoch SP, Lewis E, Hall SM, Bartlett CLR, Cohen BJ, Mortimer PP, Pereira MS. 1983. Human parvovirus, the cause of erythema infectiosum (fifth disease)? *Lancet* ii:1378.
- Breslow NE, Day DE. 1992. Statistical methods in cancer research: volume 1. The analysis of case-control studies. International Agency for Research on Cancer Scientific Publication 119. Lyon, France: IARC.
- Brown KE, Anderson SM, Young NS. 1993. Erythrocyte P antigen: cellular receptor for B19 parvovirus. *Science* 262:114–117.
- Brown KE, Young NS, Liu JM. 1994. Molecular, cellular and clinical aspects of parvovirus B19 infection. *Crit Rev Oncol Hematol* 16: 1–31.
- Brown T, Anand A, Ritchie LD, Clewley JP, Reid TMS. 1984. Intra-uterine parvovirus infection associated with hydrops fetalis. *Lancet* 2:1033–1034.
- Cassinotti P, Burtonboy G, Fopp M, Siegl G. 1997. Evidence for persistence of human parvovirus B19 DNA in bone marrow. *J Med Virol* 53:229–232.
- Cohen BJ, Buckley MM. 1988. The prevalence of antibody to human parvovirus B19 in England and Wales. *J Med Microbiol* 25:151–153.
- Cossart YE, Cant B, Field AM, Widdows D. 1975. Parvovirus-like particles in human serum. *Lancet* 1:72–73.
- Courouce AM, Ferchal F, Morinet F, Muller A, Drouet J, Soulier JP, Perol Y. 1984. Human parvovirus infection in France. *Lancet* 1: 160.
- Edwards JMB, Kessel I, Gardner SD, Eaton BR, Pollock TM, Fleck DG, Gibson P, Woodroff M, Porter AD. 1981. A search for a characteristic illness in children with serological evidence of viral or toxoplasma infection: Epstein-Barr virus, hepatitis A virus, human polyomaviruses BK and JC, parvovirus-like B₁₉ and toxoplasma. *J Infect* 3:316–323.
- Erdman DD, Anderson BC, Torok TJ, Finkel TH, Anderson LJ. 1997. Possible transmission of parvovirus B19 from intravenous immune globulin. *J Med Virol* 53:233–236.
- Gray A, Guillou L, Zufferey J, Rey F, Kurt AM, Jichlinski P, Leisinger HJ, Benhattar J. 1998. Persistence of parvovirus B19 DNA in testis of patients with testicular germ cell tumors. *J Gen Virol* 79:573–579.
- Knott PD, Welby GAC, Anderson MJ. 1984. Serologically proved intrauterine infection with parvovirus. *Br Med J* 289:1660.
- Kurtzman GJ, Cohen BJ, Field AM, Oseas R, Blacse RM, Young NS. 1989. Immune response to B19 parvovirus and an antibody defect in persistent viral infection. *J Clin Invest* 84:1114–1123.
- Lim WL, Wong KF, Lau CS. 1997. Parvovirus B19 infection in Hong Kong. *J Infect* 35:247–249.
- Matsunaga Y, Takeda N, Yamazaki S, Kamata K, Kurosana D. 1995. Seroepidemiology of human parvovirus B 19 using recombinant VP1 + VP2 particle antigen [Japanese]. *Kansenshogaku Zasshi* 69:1371–1375.
- Ministry of the Interior of R.O.C. 1985. Taiwan-Fukien demographic fact book. Taipei, Republic of China: Ministry of the Interior of Republic of China.
- Mortimer PP, Luban NLC, Kelleher JF, Cohen BJ. 1983. Transmission of serum parvovirus-like virus by clotting factor concentrates. *Lancet* ii:482–484.
- Musiani M, Zerbini M, Gentilomi G, Rodorigo G, Rosa VD, Gibellini D, Venturoli S. 1995. Persistent B19 parvovirus infections in hemophilic HIV-1 infected patients. *J Med Virol* 46:103–108.
- Nunoue T, Okochi K, Mortimer PP, Cohen BJ. 1985. Human parvovirus (B19) and erythema infectiosum. *J Pediatr* 107:38–40.
- Okabe N, Kobayashi S, Tatsuzawa O, Mortimer PP. 1984. Detection of antibodies to human parvovirus in erythema infectiosum (fifth disease). *Arch Dis Childhood* 59:1016–1019.
- Pattison JR. 1994. Human parvovirus B19. *Lancet* 308:149–150.
- Pattison JR, Jones SE, Hodgson J, Davis LR, White JM, Stroud CE, Murtaza L. 1981. Parvovirus infection and hydroplastic crisis in sickle cell anemia. *Lancet* 1:664–665.
- Paver WK, Clarke SKR. 1976. Comparison of human fecal and serum parvo-like viruses. *J Clin Microbiol* 4:67–70.
- Rao KRP, Patel AR, Anderson MJ, Hodgson J, Jones SE, Pattison JR. 1983. Infection with parvovirus-like virus and aplastic crisis in chronic hemolytic anemia. *Ann Intern Med* 98:930–932.
- Reid DM, Reid TMS, Brown T, Rennie JAN, Eastmond CJ. 1985.

- Human parvovirus associated with arthritis: a clinical and laboratory description. *Lancet* 1:422–425.
- Schwarz TF, Roggerdorf M, Deinhardt F. 1987. Prevalence of parvovirus B19 infection: a seroepidemiological study. *Deutsch Medizin Wochenschr* 112:1526–1531.
- Serjeant GR, Topley JM, Mason K, Sergeant BE, Pattison JR, Jones SE, Mohamed R. 1981. Outbreak of aplastic crises in sickle cell anemia associated with parvovirus-like agent. *Lancet* 2:595–597.
- Siegl GR, Bates RC, Bern KI, Carter BJ, Kelly DC, Kurstak E, Tattersall P. 1985. Characteristics and taxonomy of *Parvoviridae*. *Intervirology* 23:61–73.
- Shade RO, Blundell MC, Cotmore SF, Tattersall P, Astell CR. 1986. Nucleotide sequence and genome organization of human parvovirus B19 isolated from the serum of a child during aplastic crisis. *J Virol* 58:921–936.
- von Pöblitzki A, Gigler A, Lang B, Wolf H, Modrow S. 1995. Antibodies to parvovirus B19 NS-1 protein in infected individual. *J Gen Virol* 76:519–527.
- Wang CH, Chen CJ, Hu CY, You SL, Chu CT, Chou MJ, Essex M, Blattner WA, Liu CH, Yang CS. 1988. Seroepidemiology of human T-cell lymphotropic virus type I infection in Taiwan. *Cancer Res* 48:5042–5044.
- White DG, Woolf AD, Mortimer PP, Cohen BJ, Blake DR, Bacon PA. 1985. Human parvovirus arthropathy. *Lancet* 1:419–421.
- Williams MD, Cohen B, Beddal AC, Pasi K, Mortimer PP, Hill FGH. 1990. Transmission of human parvovirus B19 by coagulation factor concentrates. *Vox Sanguinis* 58:177–181.
- Yamashita K, Matsunaga Y, Taylor-Wiedeman J, Yamazaki S. 1992. A significant age shift of the human parvovirus B19 antibody prevalence among young adults in Japan observed in a decade. *Jap J Med Sci Biol* 45:49–58.